arXiv:1203.0771v1 [cond-mat.stat-mech] 4 Mar 2012

Free energy of a charged oscillator in a magnetic field and coupled to a heat bath through the momentum variables

Shamik Gupta[1] and Malay Bandyopadhyay[2]
[1]*Laboratoire de Physique de l'École Normale Supérieure de Lyon,*
Université de Lyon, CNRS, 46 Allée d'Italie, 69364 Lyon cédex 07, France
[2]*Chemical Physics Theory Group, University of Toronto,*
80, Saint George Street, Ontario M5S 3H6, Canada

(Dated: February 24, 2013)

We obtain an exact formula for the equilibrium free energy of a charged quantum particle moving in a harmonic potential in the presence of a uniform external magnetic field and linearly coupled to a heat bath through the momentum variables. The heat bath is modelled as a collection of independent quantum harmonic oscillators. The derived free energy has important differences from that in the case of coordinate-coordinate coupling between the particle and the heat-bath oscillators.

PACS numbers: 05.30.-d, 05.70.Ce

I. INTRODUCTION

Consider the dissipative dynamics of a charged quantum particle moving in a harmonic potential in the presence of a magnetic field and coupled to the external environment. Such a situation is frequently encountered in typical experiments designed to study the magnetic response of a charged quantum particle in the context of, e.g., Landau diamagnetism [1], quantum Hall effect [2], and high-temperature superconductivity [3].

In a recent paper [4], we analyzed the dynamics of the charged particle by regarding the environment as a quantum mechanical heat bath or reservoir, and by invoking a gauge-invariant system-plus-reservoir model. We considered the heat bath to consist of a collection of independent quantum harmonic oscillators, while its interaction with the charged particle was modelled in terms of bilinear coupling between the momentum variables of the particle and the oscillators. The magnetic field was taken to be uniform in space. In that work, we obtained the quantum Langevin equation (QLE) for the reduced dynamics of the particle, in which coupling to the bath is described solely by an operator-valued random force and a mean force characterized by a memory function. The magnetic field appears in the QLE through a quantum-generalized classical Lorentz force term, while the random force is independent of the field. These aspects are also present in the QLE for the case of particle-bath coordinate-coordinate coupling [5]. However, there are important differences, e.g., in the form of the memory function which now depends on the magnetic field, in the expressions for the symmetric correlation and unequal-time commutator of the random force, and also in the fact that the inertial term and the harmonic potential term in the QLE get renormalized by the coupling constants.

In this sequel paper, we further our study of the same system by deriving an exact formula for the free energy of the charged particle in thermal equilibrium. Our result shows important differences in the form of the free energy with respect to that for the coordinate-coordinate coupling [5]. Thus, this work, in addition to the previous one [4], underlines the subtle role played by the form of the coupling (coordinate-coordinate *vis-à-vis* momentum-momentum) in characterizing the dynamics and thermodynamics of the charged particle.

The paper is structured as follows. In the following Section, we define the system of study and write down the equations of motion for the charged particle and the heat-bath oscillators. Section III forms the main part of this paper in which we present a detailed derivation of the equilibrium free energy of the charged particle, and point out its differences from the case of coordinate-coordinate coupling.

II. SYSTEM OF STUDY AND EQUATIONS OF MOTION

The system considered in Ref. [4] comprises a charged quantum particle moving in a harmonic potential in the presence of a uniform external magnetic field, and linearly coupled to a quantum heat bath through the momentum variables. The heat bath is modelled as a collection of a large number of independent quantum harmonic oscillators. The Hamiltonian of the system is

$$H_0 = \frac{1}{2m}\Big(\mathbf{p} - \frac{e}{c}\mathbf{A}\Big)^2 + \frac{1}{2}m\omega_0^2\mathbf{r}^2 + \sum_{j=1}^{N}\Big[\frac{1}{2m_j}\Big(\mathbf{p}_j - g_j\mathbf{p} + \frac{g_j e}{c}\mathbf{A}\Big)^2 + \frac{1}{2}m_j\omega_j^2\mathbf{q}_j^2\Big], \tag{1}$$

where $e, m, \mathbf{p}, \mathbf{r}$ are, respectively, the charge, the mass, the momentum operator and the coordinate operator of the particle, while ω_0 is the frequency characterizing its motion in the harmonic potential. The jth heat-bath oscillator has mass m_j, frequency ω_j, coordinate operator $\mathbf{q}_j$, and momentum operator $\mathbf{p}_j$. The dimensionless parameter g_j describes the coupling between the particle and the jth oscillator. The speed of light in vacuum is denoted by c. The vector potential $\mathbf{A} = \mathbf{A}(\mathbf{r})$ is related to the uniform external magnetic field $\mathbf{B} = (B_x, B_y, B_z)$ through $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$. The field has the magnitude $B = \sqrt{B_x^2 + B_y^2 + B_z^2}$.

The commutation relations for the different coordinate and momentum operators are

$$[r_\alpha, p_\beta] = i\hbar\delta_{\alpha\beta}, [q_{j\alpha}, p_{k\beta}] = i\hbar\delta_{jk}\delta_{\alpha\beta}, \tag{2}$$

while all other commutators vanish. In the above equation, δ_{jk} denotes the Kronecker δ function. Throughout this paper, Greek indices $(\alpha, \beta, \ldots)$ refer to the three spatial directions, while Roman indices $(i, j, k, \ldots)$ represent the heat-bath oscillators. Moreover, we use Einstein summation convention for the Greek indices.

The Heisenberg equations of motion for the charged particle are

$$\begin{aligned}\dot{\mathbf{r}} &= \frac{1}{i\hbar}[\mathbf{r}, H] \\ &= \frac{1}{m}\Big(\mathbf{p} - \frac{e}{c}\mathbf{A}\Big) - \sum_{j=1}^{N}\frac{g_j}{m_j}\Big(\mathbf{p}_j - g_j\mathbf{p} + \frac{g_j e}{c}\mathbf{A}\Big),\end{aligned} \tag{3}$$

and

$$\begin{aligned}\dot{\mathbf{p}} &= \frac{1}{i\hbar}[\mathbf{p}, H] \\ &= \frac{e}{c}(\dot{\mathbf{r}} \times \mathbf{B}) + \frac{e}{c}(\dot{\mathbf{r}} \cdot \nabla)\mathbf{A} + \frac{i\hbar e}{2m_{\mathrm{r}}c}\nabla(\nabla \cdot \mathbf{A}) - m\omega_0^2\mathbf{r},\end{aligned} \tag{4}$$

where m_{r} is the "renormalized mass",

$$m_{\mathrm{r}} \equiv m/\Big[1 + \sum_{j=1}^{N}\frac{g_j^2 m}{m_j}\Big], \tag{5}$$

and dots denote differentiation with respect to time. The Heisenberg equations for the heat-bath oscillators are

$$\begin{aligned}\dot{\mathbf{q}}_j &= \frac{1}{i\hbar}[\mathbf{q}_j, H] \\ &= \frac{1}{m_j}\Big(\mathbf{p}_j - g_j\mathbf{p} + \frac{g_j e}{c}\mathbf{A}\Big),\end{aligned} \tag{6}$$

and

$$\begin{aligned}\dot{\mathbf{p}}_j &= \frac{1}{i\hbar}[\mathbf{p}_j, H] \\ &= -m_j\omega_j^2\mathbf{q}_j.\end{aligned} \tag{7}$$

These equations lead to

$$m_{\mathrm{r}}\ddot{\mathbf{r}} = -m\omega_0^2\mathbf{r} + \frac{e}{c}(\dot{\mathbf{r}} \times \mathbf{B}) + \sum_{j=1}^{N} g_j m_{\mathrm{r}}\omega_j^2\mathbf{q}_j, \tag{8}$$

and

$$m_j\ddot{\mathbf{q}}_j = -m_j\omega_j^2\mathbf{q}_j + g_j m\omega_0^2\mathbf{r} - \frac{g_j e}{c}(\dot{\mathbf{r}} \times \mathbf{B}). \tag{9}$$

For details on the derivation of Eqs. (3) - (9), see [4].

III. EQUILIBRIUM FREE ENERGY

In order to obtain the equilibrium free energy of the charged particle, we start with computing the mean energy $\langle H_0 \rangle$ of the system (1) in thermal equilibrium at temperature T. Using Eqs. (3) and (6), we get

$$\begin{aligned}\langle H_0 \rangle = &\Big[\frac{1}{2}m\langle\dot{\mathbf{r}}^2\rangle + \frac{1}{2}m\omega_0^2\langle\mathbf{r}^2\rangle\Big] \\ &+ \frac{m}{2}\Big[\sum_{j=1}^{N} g_j\langle\dot{\mathbf{r}} \cdot \dot{\mathbf{q}}_j + \dot{\mathbf{q}}_j \cdot \dot{\mathbf{r}}\rangle + \sum_{j,k=1}^{N} g_j g_k\langle\dot{\mathbf{q}}_j \cdot \dot{\mathbf{q}}_k\rangle\Big] \\ &+ \sum_{j=1}^{N}\Big[\frac{1}{2}m_j\langle\dot{\mathbf{q}}_j^2\rangle + \frac{1}{2}m_j\omega_j^2\langle\mathbf{q}_j^2\rangle\Big].\end{aligned} \tag{10}$$

Thus, to find $\langle H_0 \rangle$, we need equilibrium averages, such as $\langle\mathbf{r}^2\rangle$ and $\langle\dot{\mathbf{r}}^2\rangle$, which may be obtained by knowing the equilibrium autocorrelation function of the position of the charged particle, defined as

$$\psi_{\rho\sigma}^{rr}(t - t') \equiv \frac{1}{2}\langle r_\rho(t)r_\sigma(t') + r_\sigma(t')r_\rho(t)\rangle. \tag{11}$$

Similarly, $\langle\mathbf{q}_j^2\rangle$, $\langle\dot{\mathbf{q}}_j^2\rangle$, and $\langle\dot{\mathbf{q}}_j \cdot \dot{\mathbf{q}}_k\rangle$ are obtained from the position autocorrelation function of the heat-bath oscillators, defined as

$$\psi_{jk,\rho\sigma}^{qq}(t - t') \equiv \frac{1}{2}\langle q_{j\rho}(t)q_{k\sigma}(t') + q_{k\sigma}(t')q_{j\rho}(t)\rangle, \tag{12}$$

while $\langle\dot{\mathbf{r}}\cdot\dot{\mathbf{q}}_j + \dot{\mathbf{q}}_j\cdot\dot{\mathbf{r}}\rangle$ may be computed from the correlations

$$\psi_{j,\rho\sigma}^{rq}(t - t') \equiv \frac{1}{2}\langle r_\rho(t)q_{j\sigma}(t') + q_{j\sigma}(t')r_\rho(t)\rangle, \tag{13}$$

and

$$\psi_{j,\rho\sigma}^{qr}(t - t') \equiv \frac{1}{2}\langle q_{j,\rho}(t)r_\sigma(t') + r_\sigma(t')q_{j,\rho}(t)\rangle. \tag{14}$$

Correlations such as in Eqs. (11) - (14) are computed conveniently by invoking the fluctuation-dissipation theorem that relates such equilibrium correlations to response of the system to small external perturbations [6]. To this end, consider a weak external force $\mathbf{f}(t)$ to act on the charged particle and another set of weak external forces $\{\mathbf{f}_j(t), j = 1, 2, \ldots, N\}$ to act on the oscillators. Here, $\mathbf{f}(t)$ and $\mathbf{f}_j(t)$ are c-number functions of time. We take the perturbed Hamiltonian to be of the form

$$H = H_0 - \mathbf{r} \cdot \mathbf{f}(t) - \sum_{j=1}^{N}\mathbf{q}_j \cdot \mathbf{f}_j(t). \tag{15}$$

In the presence of these external forces, the Heisenberg equations of motion (3) and (6) remain the same, while those given by Eqs. (4) and (7), respectively, are modified to

$$\dot{\mathbf{p}} = \frac{e}{c}(\dot{\mathbf{r}} \times \mathbf{B}) + \frac{e}{c}(\dot{\mathbf{r}} \cdot \nabla)\mathbf{A} + \frac{i\hbar e}{2m_{\mathrm{r}}c}\nabla(\nabla \cdot \mathbf{A}) - m\omega_0^2\mathbf{r} + \mathbf{f}, \tag{16}$$

and

$$\dot{\mathbf{p}}_j = -m_j\omega_j^2\mathbf{q}_j + \mathbf{f}_j. \tag{17}$$

Consequently, Eqs. (8) and (9), respectively, are modified to

$$m_{\rm r}\ddot{\mathbf{r}} = -m\omega_0^2\mathbf{r} + \frac{e}{c}(\dot{\mathbf{r}}\times\mathbf{B}) + \sum_{j=1}^{N} g_j m_{\rm r}\omega_j^2\mathbf{q}_j - \sum_{j=1}^{N}\frac{g_j m_{\rm r}}{m_j}\mathbf{f}_j + \mathbf{f}, \tag{18}$$

and

$$m_j\ddot{\mathbf{q}}_j = -m_j\omega_j^2\mathbf{q}_j + g_j m\omega_0^2\mathbf{r} - \frac{g_j e}{c}(\dot{\mathbf{r}}\times\mathbf{B}) + \mathbf{f}_j - g_j\mathbf{f}. \tag{19}$$

Taking the Fourier transform of Eqs. (18) and (19), we get [7]

$$\Big[\delta_{\rho\sigma}(-m_{\rm r}\omega^2 + m\omega_0^2) + \frac{i\omega e}{c}\epsilon_{\rho\sigma\eta}B_\eta\Big]\bar{r}_\sigma - \sum_{j=1}^{N} g_j m_{\rm r}\omega_j^2\bar{q}_{j\rho}$$
$$= -\sum_{j=1}^{N}\frac{g_j m_{\rm r}}{m_j}\bar{f}_{j\rho} + \bar{f}_\rho, \tag{20}$$

and

$$(-m_j\omega^2 + m_j\omega_j^2)\bar{q}_{j\rho} - \Big(g_j m\omega_0^2\delta_{\rho\sigma} + \frac{i\omega g_j e}{c}\epsilon_{\rho\sigma\eta}B_\eta\Big)\bar{r}_\sigma$$
$$= \bar{f}_{j\rho} - g_j\bar{f}_\rho, \tag{21}$$

where $\epsilon_{\rho\sigma\eta}$ is the Levi-Civita symbol.

Equations (20) and (21) give

$$D_{\rho\sigma}(\omega)\bar{r}_\sigma = G(\omega)\bar{f}_\rho + \sum_{j=1}^{N}\frac{g_j m_{\rm r}\omega^2}{m_j(\omega_j^2-\omega^2)}\bar{f}_{j\rho}, \tag{22}$$

where

$$G(\omega) = 1 - \sum_{j=1}^{N}\frac{(g_j)^2 m_{\rm r}\omega_j^2}{m_j(\omega_j^2-\omega^2)}, \tag{23}$$

$$D_{\rho\sigma}(\omega) = \lambda(\omega)\delta_{\rho\sigma} + \frac{i\omega e G(\omega)}{c}\epsilon_{\rho\sigma\eta}B_\eta, \tag{24}$$

and

$$\lambda(\omega) = -m_{\rm r}\omega^2 + m\omega_0^2 G(\omega). \tag{25}$$

From Eq. (22), we get

$$\bar{r}_\rho = G(\omega)\alpha_{\rho\gamma}(\omega)\bar{f}_\gamma + \sum_{j=1}^{N}\beta_{j,\rho\gamma}(\omega)\bar{f}_{j\gamma}, \tag{26}$$

where

$$\alpha_{\rho\gamma}(\omega) = [D(\omega)^{-1}]_{\rho\gamma}$$
$$= \Big[(\lambda(\omega))^2\delta_{\rho\gamma} - \Big(\frac{\omega e G(\omega)}{c}\Big)^2 B_\rho B_\gamma$$
$$-\frac{i\omega\lambda(\omega)eG(\omega)}{c}\epsilon_{\rho\gamma\eta}B_\eta\Big]/\text{Det } D(\omega), \tag{27}$$

$$\text{Det } D(\omega) = \lambda(\omega)\Big[(\lambda(\omega))^2 - \Big(\frac{\omega B e G(\omega)}{c}\Big)^2\Big], \tag{28}$$

and

$$\beta_{j,\rho\gamma}(\omega) = \frac{g_j m_{\rm r}\omega^2}{m_j(\omega_j^2-\omega^2)}\alpha_{\rho\gamma}(\omega). \tag{29}$$

Using Eq. (26) in either Eq. (20) or Eq. (21) gives

$$\bar{q}_{j\rho} = \Delta_{j,\rho\gamma}(\omega)\bar{f}_\gamma + \sum_{k=1}^{N}\gamma_{jk,\rho\gamma}(\omega)\bar{f}_{k\gamma}, \tag{30}$$

where

$$\Delta_{j,\rho\gamma}(\omega) = \frac{g_j m G(\omega)}{m_j(\omega_j^2-\omega^2)}\Big(\omega_0^2\delta_{\rho\sigma} + \frac{i\omega e}{mc}\epsilon_{\rho\sigma\eta}B_\eta\Big)\alpha_{\sigma\gamma}(\omega)$$
$$-\frac{g_j}{m_j(\omega_j^2-\omega^2)}\delta_{\rho\gamma}, \tag{31}$$

and

$$\gamma_{jk,\rho\gamma}(\omega)$$
$$= \frac{g_j g_k m m_{\rm r}\omega^2}{m_j m_k(\omega_j^2-\omega^2)(\omega_k^2-\omega^2)}\Big(\omega_0^2\delta_{\rho\sigma} + \frac{i\omega e}{mc}\epsilon_{\rho\sigma\eta}B_\eta\Big)\alpha_{\sigma\gamma}(\omega)$$
$$+\frac{\delta_{jk}\delta_{\rho\gamma}}{m_k(\omega_k^2-\omega^2)}. \tag{32}$$

The functions $\alpha_{\rho\gamma}(\omega)$, $\beta_{j,\rho\gamma}(\omega)$, $\Delta_{j,\rho\gamma}(\omega)$, and $\gamma_{jk,\rho\gamma}(\omega)$ are coefficient matrices of the response of the system to the external perturbations, $\mathbf{f}(t)$ and $\mathbf{f}_j(t)$, and may be interpreted as generalized susceptibilities.

We may now use the fluctuation-dissipation theorem which relates the Fourier transform $\bar{\psi}^{rr}_{\rho\sigma}(\omega)$ of the position autocorrelation function $\psi^{rr}_{\rho\sigma}(t-t')$ with $G(\omega)\alpha_{\rho\sigma}(\omega)$:

$$\bar{\psi}^{rr}_{\rho\sigma}(\omega) = \frac{\hbar}{2i}\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]G(\omega)[\alpha_{\rho\sigma}(\omega+i0^+)$$
$$-\alpha^*_{\sigma\rho}(\omega+i0^+)], \tag{33}$$

where $*$ denotes complex conjugation (Note that $G(\omega)$ is real). Similarly, one has

$$\bar{\psi}^{qq}_{jk,\rho\sigma}(\omega) = \frac{\hbar}{2i}\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]$$
$$\times[\gamma_{jk,\rho\sigma}(\omega+i0^+) - \gamma^*_{jk,\sigma\rho}(\omega+i0^+)], \tag{34}$$

$$\bar{\psi}^{rq}_{j,\rho\sigma}(\omega) = \frac{\hbar}{2i}\coth\Big[\frac{\hbar\omega}{2k_BT}\Big][\beta_{j,\rho\sigma}(\omega+i0^+) - \beta^*_{j,\sigma\rho}(\omega+i0^+)], \tag{35}$$

and

$$\bar{\psi}^{qr}_{j,\rho\sigma}(\omega) = \frac{\hbar}{2i}\coth\Big[\frac{\hbar\omega}{2k_BT}\Big][\Delta_{j,\rho\sigma}(\omega+i0^+) - \Delta^*_{j,\sigma\rho}(\omega+i0^+)]. \tag{36}$$

In passing, we note that

$$\alpha^*_{\sigma\rho}(\omega) = \alpha_{\sigma\rho}(-\omega), \tag{37}$$

which implies that $\mathrm{Im}[\alpha_{\sigma\rho}(\omega)]$ is an odd function of ω, while $\mathrm{Re}[\alpha_{\sigma\rho}(\omega)]$ is an even function of ω. These properties are also shared by $\beta_{j,\sigma\rho}(\omega), \Delta_{j,\sigma\rho}(\omega)$, and $\gamma_{jk,\sigma\rho}(\omega)$.

In the following, we use Eqs. (33) - (36) to evaluate the various averages involved in computing $\langle H_0\rangle$.

A. Computation of various averages

1. Averages involving $\mathbf{r}$

We start with decomposing $\alpha_{\rho\sigma}$ into symmetric and antisymmetric parts as

$$\alpha_{\rho\sigma}(\omega) = \alpha^{\mathrm{s}}_{\rho\sigma}(\omega) + \alpha^{\mathrm{a}}_{\rho\sigma}(\omega). \tag{38}$$

We then have

$$\begin{aligned}&\alpha_{\rho\sigma}(\omega) - \alpha^*_{\sigma\rho}(\omega)\\ &= [\alpha^{\mathrm{s}}_{\rho\sigma}(\omega) - \alpha^{\mathrm{s}}_{\rho\sigma}(\omega)^*] + [\alpha^{\mathrm{a}}_{\rho\sigma}(\omega) + \alpha^{\mathrm{a}}_{\rho\sigma}(\omega)^*]\\ &= 2i\mathrm{Im}[\alpha^{\mathrm{s}}_{\rho\sigma}(\omega)] + 2\mathrm{Re}[\alpha^{\mathrm{a}}_{\rho\sigma}(\omega)].\end{aligned} \tag{39}$$

Using Eqs. (11), (33), (39), and the properties that $\mathrm{Im}[\alpha_{\rho\sigma}(\omega)]$ is an odd function of ω, while both $G(\omega)$ and $\mathrm{Re}[\alpha_{\rho\sigma}(\omega)]$ are even functions of ω, we get

$$\begin{aligned}&\frac{1}{2}\langle r_\rho(t)r_\sigma(t') + r_\sigma(t')r_\rho(t)\rangle\\ &= \frac{\hbar}{\pi}\int_0^\infty d\omega\ G(\omega)\mathrm{Im}[\alpha^{\mathrm{s}}_{\rho\sigma}(\omega+i0^+)]\\ &\qquad\times\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\cos[\omega(t-t')]\\ &-\frac{\hbar}{\pi}\int_0^\infty d\omega\ G(\omega)\mathrm{Re}[\alpha^{\mathrm{a}}_{\rho\sigma}(\omega+i0^+)]\\ &\qquad\times\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\sin[\omega(t-t')],\end{aligned} \tag{40}$$

which for $\rho=\sigma$ gives

$$\begin{aligned}&\frac{1}{2}\langle \mathbf{r}(t)\cdot\mathbf{r}(t') + \mathbf{r}(t')\cdot\mathbf{r}(t)\rangle\\ &= \frac{\hbar}{\pi}\int_0^\infty d\omega\ G(\omega)\mathrm{Im}[\alpha_{\rho\rho}(\omega+i0^+)]\\ &\qquad\times\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\cos[\omega(t-t')].\end{aligned} \tag{41}$$

Then, putting $t=t'$, we get

$$\langle\mathbf{r}^2\rangle = \frac{\hbar}{\pi}\int_0^\infty d\omega\ G(\omega)\mathrm{Im}[\alpha_{\rho\rho}(\omega+i0^+)]\coth\Big[\frac{\hbar\omega}{2k_BT}\Big], \tag{42}$$

where from Eq. (27), we have

$$\alpha_{\rho\rho}(\omega) = \Big[(\lambda(\omega))^2\delta_{\rho\rho} - \Big(\frac{\omega eG(\omega)}{c}\Big)^2 B_\rho B_\rho\Big]/\mathrm{Det}\ D(\omega). \tag{43}$$

Differentiating Eq. (41) successively with respect to t and t', and finally putting $t=t'$, we get

$$\begin{aligned}&\langle\dot{\mathbf{r}}^2\rangle\\ &= \frac{\hbar}{\pi}\int_0^\infty d\omega\ G(\omega)\mathrm{Im}[\alpha_{\rho\rho}(\omega+i0^+)]\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\omega^2.\end{aligned} \tag{44}$$

2. Averages involving $\mathbf{q}_j$

One may proceed similarly to the preceding subsection to obtain

$$\begin{aligned}&\frac{1}{2}\langle\mathbf{q}_j(t)\cdot\mathbf{q}_k(t') + \mathbf{q}_k(t')\cdot\mathbf{q}_j(t)\rangle\\ &= \frac{\hbar}{\pi}\int_0^\infty d\omega\ \mathrm{Im}[\gamma_{jk,\rho\rho}(\omega+i0^+)]\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\cos[\omega(t-t')],\end{aligned} \tag{45}$$

where $\gamma_{jk,\rho\rho}(\omega)$ is obtained from Eq. (32) as

$$\begin{aligned}\gamma_{jk,\rho\rho}(\omega) &= \frac{g_jg_kmm_{\mathrm{r}}\omega^2\omega_0^2}{m_jm_k(\omega_j^2-\omega^2)(\omega_k^2-\omega^2)}\alpha_{\rho\rho}(\omega)\\ &-\frac{g_jg_km_{\mathrm{r}}\omega^4\lambda(\omega)e^2G(\omega)}{m_jm_k(\omega_j^2-\omega^2)(\omega_k^2-\omega^2)c^2\mathrm{Det}\ D(\omega)}(\delta_{\rho\rho}B^2 - B_\rho B_\rho)\\ &+\frac{\delta_{jk}\delta_{\rho\rho}}{m_k(\omega_k^2-\omega^2)}.\end{aligned} \tag{46}$$

One therefore gets

$$\langle\mathbf{q}_j^2\rangle = \frac{\hbar}{\pi}\int_0^\infty d\omega\ \mathrm{Im}[\gamma_{jj,\rho\rho}(\omega+i0^+)]\coth\Big[\frac{\hbar\omega}{2k_BT}\Big], \tag{47}$$

$$\langle\dot{\mathbf{q}}_j^2\rangle = \frac{\hbar}{\pi}\int_0^\infty d\omega\ \mathrm{Im}[\gamma_{jj,\rho\rho}(\omega+i0^+)]\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\omega^2, \tag{48}$$

and

$$\langle\dot{\mathbf{q}}_j\cdot\dot{\mathbf{q}}_k\rangle = \frac{\hbar}{\pi}\int_0^\infty d\omega\ \mathrm{Im}[\gamma_{jk,\rho\rho}(\omega+i0^+)]\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\omega^2. \tag{49}$$

3. Averages involving $\mathbf{r}$ and $\mathbf{q}_j$

We have

$$\frac{1}{2}\langle r_\rho(t)q_{j\sigma}(t')+q_{j\sigma}(t')r_\rho(t)\rangle = \frac{1}{2\pi}\int_{-\infty}^\infty d\omega\ e^{-i\omega(t-t')}\bar{\psi}^{rq}_{j,\rho\sigma}(\omega), \tag{50}$$

and

$$\frac{1}{2}\langle q_{j\sigma}(t')r_\rho(t)+r_\rho(t)q_{j\sigma}(t')\rangle = \frac{1}{2\pi}\int_{-\infty}^\infty d\omega\ e^{-i\omega(t'-t)}\bar{\psi}^{qr}_{j,\sigma\rho}(\omega). \tag{51}$$

Adding Eq. (51) to Eq. (50), we get

$$\langle r_\rho(t) q_{j\sigma}(t') + q_{j\sigma}(t') r_\rho(t)\rangle \\ = \frac{1}{2\pi}\int_{-\infty}^{\infty} d\omega\ \Big[e^{-i\omega(t-t')}\bar{\psi}^{rq}_{j,\rho\sigma}(\omega) + e^{-i\omega(t'-t)}\bar{\psi}^{qr}_{j,\sigma\rho}(\omega)\Big]. \tag{52}$$

Differentiating Eq. (52) successively with respect to t and t', and then putting $t = t'$, we get

$$\langle \dot{r}_\rho \dot{q}_{j\sigma} + \dot{q}_{j\sigma}\dot{r}_\rho\rangle = \frac{1}{2\pi}\int_{-\infty}^{\infty} d\omega\ \omega^2\Big[\bar{\psi}^{rq}_{j,\rho\sigma}(\omega) + \bar{\psi}^{qr}_{j,\sigma\rho}(\omega)\Big]. \tag{53}$$

Using Eqs. (35) and (36) for $\bar{\psi}^{rq}_{j,\rho\sigma}(\omega)$ and $\bar{\psi}^{qr}_{j,\sigma\rho}(\omega)$, respectively, and noting that the imaginary part of both $\beta_{j,\rho\sigma}(\omega)$ and $\Delta_{j,\rho\sigma}(\omega)$ is an odd function of ω, while their real parts are even functions of ω, we get

$$\langle \dot{\mathbf{q}}_j\cdot\dot{\mathbf{r}} + \dot{\mathbf{r}}\cdot\dot{\mathbf{q}}_j\rangle \\ = \frac{\hbar}{\pi}\int_0^\infty d\omega\ \coth\Big[\frac{\hbar\omega}{2k_BT}\Big] \\ \times\omega^2\Big[\mathrm{Im}[\beta_{j,\rho\rho}(\omega+i0^+)] + \mathrm{Im}[\Delta_{j,\rho\rho}(\omega+i0^+)]\Big], \tag{54}$$

where from Eq. (31), we get

$$\Delta_{j,\rho\rho}(\omega) = \frac{g_j m G(\omega)\omega_0^2}{m_j(\omega_j^2-\omega^2)}\alpha_{\rho\rho}(\omega) \\ -\frac{g_j (G(\omega))^2\lambda(\omega)\omega^2 e^2}{m_j(\omega_j^2-\omega^2)c^2\mathrm{Det}\ D(\omega)}(\delta_{\rho\rho}B^2 - B_\rho B_\rho) \\ -\frac{g_j}{m_j(\omega_j^2-\omega^2)}\delta_{\rho\rho}. \tag{55}$$

B. Mean energy

Using Eqs. (42), (44), (47), (48), (49), and (54) in Eq. (10), we get

$$\langle H_0\rangle = \frac{\hbar}{\pi}\int_0^\infty d\omega\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\Big[\frac{1}{2}m(\omega^2+\omega_0^2)G(\omega)\mathrm{Im}[\alpha_{\rho\rho}(\omega+i0^+)] \\ +\sum_{j=1}^{N}\frac{m g_j\omega^2}{2}\Big(\mathrm{Im}[\beta_{j,\rho\rho}(\omega+i0^+)] + \mathrm{Im}[\Delta_{j,\rho\rho}(\omega+i0^+)]\Big) + \sum_{j,k=1}^{N}\frac{m g_j g_k\omega^2}{2}\mathrm{Im}[\gamma_{jk,\rho\rho}(\omega+i0^+)] \\ +\sum_{j=1}^{N}\frac{1}{2}m_j(\omega^2+\omega_j^2)\mathrm{Im}[\gamma_{jj,\rho\rho}(\omega+i0^+)]\Big]. \tag{56}$$

The mean energy of the heat bath is

$$\langle H_B\rangle = \sum_{j=1}^{N}\frac{1}{2}m_j\omega_j^2\langle\mathbf{q}_j^2\rangle + \sum_{j=1}^{N}\frac{1}{2}m_j\langle\dot{\mathbf{q}}_j^2\rangle = \sum_{j=1}^{N}\frac{m_j\hbar}{2\pi}\int_0^\infty d\omega\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\mathrm{Im}[\gamma_{jj,\rho\rho}(\omega+i0^+)](\omega^2+\omega_j^2) \\ = \sum_{j=1}^{N}\frac{3\hbar}{2\pi}\int_0^\infty d\omega\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\Big(\frac{\omega^2+\omega_j^2}{\omega_j^2-\omega^2}\Big) \\ +\frac{\hbar}{2\pi}\int_0^\infty d\omega\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\sum_{j=1}^{N}m_j(\omega^2+\omega_j^2)\mathrm{Im}\Big[\frac{(g_j)^2 m m_{\mathrm{r}}\omega^2\omega_0^2}{(m_j)^2(\omega_j^2-\omega^2)^2}\alpha_{\rho\rho} - \frac{2(g_j)^2 m_{\mathrm{r}}\omega^4\lambda B^2 e^2 G}{(m_j)^2(\omega_j^2-\omega^2)^2c^2\mathrm{Det}\ D}\Big] \tag{57}$$

$$= \sum_{j=1}^{N}\frac{3\hbar\omega_j}{2}\coth\Big[\frac{\hbar\omega_j}{2k_BT}\Big] \\ +\frac{\hbar}{2\pi}\int_0^\infty d\omega\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\sum_{j=1}^{N}m_j(\omega^2+\omega_j^2)\mathrm{Im}\Big[\frac{(g_j)^2 m m_{\mathrm{r}}\omega^2\omega_0^2}{(m_j)^2(\omega_j^2-\omega^2)^2}\alpha_{\rho\rho} - \frac{2(g_j)^2 m_{\mathrm{r}}\omega^4\lambda B^2 e^2 G}{(m_j)^2(\omega_j^2-\omega^2)^2c^2\mathrm{Det}\ D}\Big]. \tag{58}$$

Here, in obtaining Eq. (57), we have used Eq. (46). To arrive at Eq. (58), we perform the first integral on the right hand side of Eq. (57) by noting that ω in the integral is approached from above the real axis, $\omega\to\omega+i0^+$, and by using the result that

$$\frac{1}{\omega-\omega_j+i0^+} = \mathrm{P}\Big[\frac{1}{\omega-\omega_j}\Big] - i\pi\delta(\omega-\omega_j), \tag{59}$$

with P denoting the principal value. Note that the first term on the right hand side of Eq. (58) is the mean energy $U_B(T)$ of the heat bath in the absence of any coupling with the charged particle:

$$U_B(T) = \sum_{j=1}^{N} \frac{3\hbar\omega_j}{2} \coth\Big[\frac{\hbar\omega_j}{2k_BT}\Big]. \tag{60}$$

We define the particle energy $U_0(T, B)$ as the mean energy of the system of the charged particle interacting with the heat bath minus the mean energy of the heat bath in the absence of coupling with the particle:

$$\begin{aligned}
&U_0(T,B) \equiv \langle H_0\rangle - U_B(T)\\
&= \frac{\hbar}{\pi}\int_0^\infty d\omega \coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\\
&\times\Big[\frac{1}{2}m(\omega^2+\omega_0^2)G\mathrm{Im}[\alpha_{\rho\rho}]\\
&+\sum_{j=1}^{N}\frac{mg_j\omega^2}{2}\Big(\mathrm{Im}\Big[\frac{g_jm_\mathrm{r}\omega^2}{m_j(\omega_j^2-\omega^2)}\alpha_{\rho\rho}\Big]+\mathrm{Im}\Big[\frac{g_jmG\omega_0^2}{m_j(\omega_j^2-\omega^2)}\alpha_{\rho\rho}-\frac{2g_jG^2\lambda\omega^2e^2B^2}{m_j(\omega_j^2-\omega^2)c^2\mathrm{Det}\ D}-\frac{3g_j}{m_j(\omega_j^2-\omega^2)}\Big]\Big)\\
&+\sum_{j,k=1}^{N}\frac{mg_jg_k\omega^2}{2}\mathrm{Im}\Big[\frac{g_jg_kmm_\mathrm{r}\omega^2\omega_0^2}{m_jm_k(\omega_j^2-\omega^2)(\omega_k^2-\omega^2)}\alpha_{\rho\rho}-\frac{2g_jg_km_\mathrm{r}\omega^4\lambda e^2GB^2}{m_jm_k(\omega_j^2-\omega^2)(\omega_k^2-\omega^2)c^2\mathrm{Det}\ D}+\frac{3\delta_{jk}}{m_k(\omega_k^2-\omega^2)}\Big]\\
&+\sum_{j=1}^{N}\frac{1}{2}m_j(\omega_j^2+\omega^2)\mathrm{Im}\Big[\frac{(g_j)^2mm_\mathrm{r}\omega^2\omega_0^2}{(m_j)^2(\omega_j^2-\omega^2)^2}\alpha_{\rho\rho}-\frac{2(g_j)^2m_\mathrm{r}\omega^4\lambda B^2e^2G}{(m_j)^2(\omega_j^2-\omega^2)^2c^2\mathrm{Det}\ D}\Big]\Big],
\end{aligned} \tag{61}$$

where we have used Eqs (29), (46), and (55). After simplification, we get

$$\begin{aligned}
&U_0(T,B)\\
&=\frac{\hbar}{\pi}\int_0^\infty d\omega\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]\mathrm{Im}\Big[\Big\{\frac{1}{2}m_\mathrm{r}(\omega^2+\omega_0^2)+\sum_{j=1}^{N}\frac{(g_j)^2mm_\mathrm{r}\omega^2\omega_0^2(\omega_j^2+\omega^2)}{2m_j(\omega_j^2-\omega^2)^2}\Big\}\alpha_{\rho\rho}+\lambda\Big(\frac{\omega eB}{c}\Big)^2\Big(\frac{dG^2}{d\omega}\Big)\frac{\omega}{2\mathrm{Det}\ D}\Big].
\end{aligned} \tag{62}$$

From Eqs. (27) and (28), we have

$$\begin{aligned}
\mathrm{Det}\ \alpha(\omega) &= [\mathrm{Det}\ D(\omega)]^{-1}\\
&= \Big[\lambda(\omega)\Big[(\lambda(\omega))^2-\Big(\frac{\omega BeG(\omega)}{c}\Big)^2\Big]\Big]^{-1},
\end{aligned} \tag{63}$$

while the trace of $\alpha(\omega)$ is

$$\alpha_{\rho\rho}(\omega) = \Big[3(\lambda(\omega))^2-\Big(\frac{\omega eG(\omega)}{c}\Big)^2B^2\Big]/\mathrm{Det}\ D(\omega). \tag{64}$$

Following [5], we write

$$\begin{aligned}
&\omega\frac{d}{d\omega}\ln[\mathrm{Det}\ \alpha(\omega)]\\
&= -\omega\Big\{\frac{d\lambda(\omega)}{d\omega}\Big[3(\lambda(\omega))^2-\Big(\frac{\omega BeG(\omega)}{c}\Big)^2\Big]\\
&\qquad -2\omega\lambda(\omega)\Big(\frac{BeG(\omega)}{c}\Big)^2\Big\}/\mathrm{Det}\ D(\omega)\\
&= -3+\Big[\lambda(\omega)-\omega\frac{d\lambda(\omega)}{d\omega}\Big]\alpha_{\rho\rho}(\omega).
\end{aligned} \tag{65}$$

From Eq. (25), we get

$$\begin{aligned}
&\lambda(\omega)-\omega\frac{d\lambda(\omega)}{d\omega}\\
&= m_\mathrm{r}(\omega^2+\omega_0^2)+\sum_{j=1}^{N}\frac{(g_j)^2mm_\mathrm{r}\omega^2\omega_0^2(\omega_j^2+\omega^2)}{m_j(\omega_j^2-\omega^2)^2}.
\end{aligned} \tag{66}$$

Now, using Eqs. (63), (65) and (66) in Eq. (62), we get

$$\begin{aligned}
U_0(T,B) &= \frac{1}{\pi}\int_0^\infty d\omega\ u(\omega,T)\\
&\times\mathrm{Im}\Big[\frac{d}{d\omega}\ln[\mathrm{Det}\ \alpha(\omega+i0^+)]\\
&\qquad +\lambda\Big(\frac{\omega eB}{c}\Big)^2\Big(\frac{dG^2}{d\omega}\Big)\mathrm{Det}\ \alpha(\omega+i0^+)\Big],
\end{aligned} \tag{67}$$

where $u(\omega, T)$ is the Planck energy of a free oscillator of frequency ω:

$$u(\omega,T) = \frac{\hbar\omega}{2}\coth\Big[\frac{\hbar\omega}{2k_BT}\Big]. \tag{68}$$

C. Free energy

The free energy $F_0(T,B)$ of the charged particle is obtained from Eq. (67), since the two quantities are related as $U_0(T,B) = F_0(T,B) - T\frac{\partial F_0(T,B)}{\partial T}$. We get

$$\begin{aligned} F_0(T,B) &= \frac{1}{\pi}\int_0^\infty d\omega\ f(\omega,T) \\ &\times \mathrm{Im}\Big[\frac{d}{d\omega}\ln[\mathrm{Det}\ \alpha(\omega+i0^+)] \\ &+\lambda\Big(\frac{\omega eB}{c}\Big)^2\Big(\frac{dG^2}{d\omega}\Big)\mathrm{Det}\ \alpha(\omega+i0^+)\Big], \end{aligned} \tag{69}$$

where $f(\omega,T)$ is the free energy of a free oscillator of frequency ω:

$$f(\omega,T) = k_BT\ln\Big[2\sinh\Big[\frac{\hbar\omega}{2k_BT}\Big]\Big]. \tag{70}$$

Equation (69) is the central result of the paper. To make explicit the contribution of the external magnetic field to the free energy, we write Eq. (63) as

$$\mathrm{Det}\ \alpha(\omega) = [\alpha^{(0)}(\omega)]^3\Big[1-\Big(\frac{\omega BeG(\omega)}{c}\Big)^2[\alpha^{(0)}(\omega)]^2\Big]^{-1}, \tag{71}$$

where

$$\alpha^{(0)}(\omega) = 1/\lambda(\omega) \tag{72}$$

is the susceptibility in the absence of the magnetic field. Using Eq. (71) in Eq. (69), we get

$$F_0(T,B) = F_0(T,0) + \Delta_1F_0(T,B) + \Delta_2F_0(T,B), \tag{73}$$

where

$$F_0(T,0) = \frac{3}{\pi}\int_0^\infty d\omega\ f(\omega,T)\mathrm{Im}\Big[\frac{d}{d\omega}\ln\alpha^{(0)}(\omega+i0^+)\Big] \tag{74}$$

is the free energy of the charged particle in the absence of the magnetic field. The contribution from the latter is contained in the two terms $\Delta_1F_0(T,B)$ and $\Delta_2F_0(T,B)$, given by

$$\begin{aligned} \Delta_1F_0(T,B) &= -\frac{1}{\pi}\int_0^\infty d\omega\ f(\omega,T) \\ &\times\mathrm{Im}\Big[\frac{d}{d\omega}\ln\Big\{1-G^2\Big(\frac{\omega Be}{c}\Big)^2[\alpha^{(0)}(\omega+i0^+)]^2\Big\}\Big], \end{aligned} \tag{75}$$

and

$$\begin{aligned} \Delta_2F_0(T,B) &= \frac{1}{\pi}\int_0^\infty d\omega\ f(\omega,T) \\ &\mathrm{Im}\Big[[\alpha^{(0)}(\omega+i0^+)]^2\Big(\frac{\omega eB}{c}\Big)^2 \\ &\Big(\frac{dG^2}{d\omega}\Big)\Big\{1-\Big(\frac{\omega BeG}{c}\Big)^2[\alpha^{(0)}(\omega+i0^+)]^2\Big\}^{-1}\Big]. \end{aligned} \tag{76}$$

Let us now comment on the form of the free energy (73) with respect to that for coordinate-coordinate coupling between the particle and the heat-bath oscillators, obtained in Ref. [5]. In the latter case, the free energy is given by

$$F_0(T,B) = F_0(T,0) + \Delta_1F_0(T,B), \tag{77}$$

where $F_0(T,0)$ has the same form as in Eq. (74), while $\Delta_1F_0(T,B)$ is given by

$$\begin{aligned} \Delta_1F_0(T,B) &= -\frac{1}{\pi}\int_0^\infty d\omega\ f(\omega,T) \\ &\times\mathrm{Im}\Big[\frac{d}{d\omega}\ln\Big\{1-\Big(\frac{\omega Be}{c}\Big)^2[\alpha^{(0)}(\omega+i0^+)]^2\Big\}\Big]. \end{aligned} \tag{78}$$

We thus see that with respect to coordinate-coordinate coupling, the free energy has two differences, namely, (i) the appearance of the extra factor G^2 in the term $\Delta_1F_0(T,B)$, and (ii) the presence of the additional term $\Delta_2F_0(T,B)$. It is interesting to see that in the absence of the magnetic field, the free energy has the same expression irrespective of the form of coupling.

With respect to the two schemes of coupling, the differences observed here in the form of the thermodynamic function, i.e., the free energy, add to the ones noted in the reduced dynamical description of the charged particle by means of the quantum Langevin equation [4]. This work thus re-emphasizes the crucial role played by the form of the coupling with the heat bath in characterizing the behavior of the charged particle. Investigations to obtain the different thermodynamic properties of the charged particle, e.g., its entropy and specific heat, by using Eq. (73) are in progress.

IV. ACKNOWLEDGMENTS

S. G. thanks the Weizmann Institute of Science, Israel, where this work was initiated. He acknowledges support of the Israel Science Foundation and the French contract ANR-10-CEXC-010-01. M. B. acknowledges support of the Connaught fund and the NSERC.

---

[1] J.H. Van Vleck, *The Theory of Electric and Magnetic Susceptibilities* (Oxford University Press, London, 1932).

[2] R.B. Laughlin, Phys. Rev. B **23**, 5632 (1981).

[3] V.L. Ginzburg and D.A. Kirzhntis, *High Temperature Superconductivity* (Consultants Bureau, NY, 1982).

[4] S. Gupta and M. Bandyopadhyay, Phys. Rev. E **84**, 041133 (2011).

[5] X.L. Li, G.W. Ford, and R.F. O'Connell, Phys. Rev. A **42**, 4519 (1990).

[6] R. Kubo, Rep. Prog. Phys. **29**, 255 (1966).

[7] For any function $g(t)$, we take its Fourier transform $\bar{g}$ to be given by

$$\bar{g}(\omega) = \int_{-\infty}^{\infty} dt\ g(t)e^{i\omega t},$$

so that

$$g(t) = \frac{1}{2\pi}\int_{-\infty}^{\infty} d\omega\ \bar{g}(\omega)e^{-i\omega t}.$$